

【 参考文献 】

I 引用文献

- (1) メタボリックシンドローム診断基準検討委員会. メタボリックシンドロームの定義と診断基準. 日本内科学雑誌 2005;94:188-203.
- (2) Carroll S, Cooke CB, Butterly RJ. Physical activity, cardiorespiratory fitness, and the primary components of blood viscosity. Med Sci Sports Exerc 2000;32:353-8.
- (3) Dipietro L, Caspersen CJ, Ostfeld AM, Nadel ER. A survey for assessing physical activity among older adults. Med Sci Sports Exerc 1993;25:628-42.
- (4) Nagaya T, Kondo Y, Shibata T. Effects of sedentary work on physical fitness and serum cholesterol profile in middle-aged male workers. Int Arch Occup Environ Health 2001;74:366-70.
- (5) Huang G, Gibson CA, Tran ZV, Osness WH. Controlled endurance exercise training and VO_{2max} changes in older adults: a meta-analysis. Prev Cardiol 2005;8:217-25.
- (6) Rauramaa R, Tuomainen P, Vaisanen S, Rankinen T. Physical activity and health-related fitness in middle-aged men. Med Sci Sports Exerc 1995;27:707-12.
- (7) Leon AS, Casal D, Jacobs D, Jr. Effects of 2,000 kcal per week of walking and stair climbing on physical fitness and risk factors for coronary heart disease. J Cardiopulm Rehabil 1996;16:183-92.
- (8) Wolfarth B, Bray MS, Hagberg JM, et al. The human gene map for performance and health-related fitness phenotypes: the 2004 update. Med Sci Sports Exerc 2005;37:881-903.
- (9) Talbot LA, Morrell CH, Metter EJ, Fleg JL. Comparison of cardiorespiratory fitness versus leisure time physical activity as predictors of coronary events in men aged < or = 65 years and > 65 years. Am J Cardiol 2002;89:1187-92.
- (10) Pate RR, Pratt M, Blair SN, et al. Physical activity and public health. A recommendation from the Centers for Disease Control and Prevention and the American College of Sports Medicine. Jama 1995;273:402-7.
- (11) Washburn RA, Seals DR. Peak oxygen uptake during arm cranking for men and women. J Appl Physiol 1984;56:954-7.
- (12) Hermansen L, Saltin B. Oxygen uptake during maximal treadmill and bicycle exercise. J Appl Physiol 1969;26:31-7.

II 上記以外に運動基準の策定に用いた文献

1. 健康の維持・増進に必要な身体活動量決定に参考とした文献

1. Salonen JT, Puska P, Tuomilehto J. Physical activity and risk of myocardial infarction, cerebral stroke and death: a longitudinal study in Eastern Finland. *Am J Epidemiol* 1982;115:526-37.
2. Paffenbarger RS, Jr., Wing AL, Hyde RT, Jung DL. Physical activity and incidence of hypertension in college alumni. *Am J Epidemiol* 1983;117:245-57.
3. Paffenbarger RS, Jr., Hyde RT, Wing AL, Hsieh CC. Physical activity, all-cause mortality, and longevity of college alumni. *N Engl J Med* 1986;314:605-13.
4. Slattery ML, Jacobs DR, Jr., Nichaman MZ. Leisure time physical activity and coronary heart disease death. The US Railroad Study. *Circulation* 1989;79:304-11.
5. Helmrich SP, Ragland DR, Leung RW, Paffenbarger RS, Jr. Physical activity and reduced occurrence of non-insulin-dependent diabetes mellitus. *N Engl J Med* 1991;325:147-52.
6. Manson JE, Rimm EB, Stampfer MJ, et al. Physical activity and incidence of non-insulin-dependent diabetes mellitus in women. *Lancet* 1991;338:774-8.
7. Manson JE, Nathan DM, Krolewski AS, Stampfer MJ, Willett WC, Hennekens CH. A prospective study of exercise and incidence of diabetes among US male physicians. *Jama* 1992;268:63-7.
8. Wannamethee G, Shaper AG. Physical activity and stroke in British middle aged men. *Bmj* 1992;304:597-601.
9. Lindenstrom E, Boysen G, Nyboe J. Lifestyle factors and risk of cerebrovascular disease in women. The Copenhagen City Heart Study. *Stroke* 1993;24:1468-72.
10. Lindenstrom E, Boysen G, Nyboe J. Risk factors for stroke in Copenhagen, Denmark. II. Life-style factors. *Neuroepidemiology* 1993;12:43-50.
11. Paffenbarger RS, Jr., Kampert JB, Lee IM, Hyde RT, Leung RW, Wing AL. Changes in physical activity and other lifeway patterns influencing longevity. *Med Sci Sports Exerc* 1994;26:857-65.
12. Lee IM, Hsieh CC, Paffenbarger RS, Jr. Exercise intensity and longevity in men. The Harvard Alumni Health Study. *Jama* 1995;273:1179-84.
13. Ching PL, Willett WC, Rimm EB, Colditz GA, Gortmaker SL, Stampfer MJ. Activity level and risk of overweight in male health professionals. *Am J Public Health* 1996;86:25-30.
14. Lynch J, Helmrich SP, Lakka TA, et al. Moderately intense physical activities and high levels of cardiorespiratory fitness reduce the risk of non-insulin-dependent diabetes mellitus in middle-aged men. *Arch Intern Med* 1996;156:1307-14.

15. 森岡聖次. コホート研究による生命予後に影響を及ぼす日常生活習慣要因の検討. 日本公衆衛生雑誌 1996;43:469-478.
16. Haapanen N, Miilunpalo S, Vuori I, Oja P, Pasanen M. Association of leisure time physical activity with the risk of coronary heart disease, hypertension and diabetes in middle-aged men and women. Int J Epidemiol 1997;26:739-47.
17. Kushi LH, Fee RM, Folsom AR, Mink PJ, Anderson KE, Sellers TA. Physical activity and mortality in postmenopausal women. JAMA 1997;277:1287-92.
18. 吉岡輝彦 岩, 大城等. 農山村における生活習慣に関するコホート研究 生活習慣と死亡との関係. 米子医学雑誌 1997;48:164-170.
19. Joakimsen RM, Fonnebo V, Magnus JH, Stormer J, Tolland A, Sogaard AJ. The Tromso Study: physical activity and the incidence of fractures in a middle-aged population. J Bone Miner Res 1998;13:1149-57.
20. Kujala UM, Kaprio J, Sarna S, Koskenvuo M. Relationship of leisure-time physical activity and mortality: the Finnish twin cohort. JAMA 1998;279:440-4.
21. Lee IM, Paffenbarger RS, Jr. Physical activity and stroke incidence: the Harvard Alumni Health Study. Stroke 1998;29:2049-54.
22. Hayashi T, Tsumura K, Suematsu C, Okada K, Fujii S, Endo G. Walking to work and the risk for hypertension in men: the Osaka Health Survey. Ann Intern Med 1999;131:21-6.
23. Hu FB, Sigal RJ, Rich-Edwards JW, et al. Walking compared with vigorous physical activity and risk of type 2 diabetes in women: a prospective study. JAMA 1999;282:1433-9.
24. Hu FB, Stampfer MJ, Colditz GA, et al. Physical activity and risk of stroke in women. JAMA 2000;283:2961-7.
25. Kujala UM, Kaprio J, Kannus P, Sarna S, Koskenvuo M. Physical activity and osteoporotic hip fracture risk in men. Arch Intern Med 2000;160:705-8.
26. Okada K, Hayashi T, Tsumura K, Suematsu C, Endo G, Fujii S. Leisure-time physical activity at weekends and the risk of Type 2 diabetes mellitus in Japanese men: the Osaka Health Survey. Diabet Med 2000;17:53-8.
27. Hoidrup S, Sorensen TI, Stroger U, Lauritzen JB, Schroll M, Gronbaek M. Leisure-time physical activity levels and changes in relation to risk of hip fracture in men and women. Am J Epidemiol 2001;154:60-8.
28. Feskanich D, Willett W, Colditz G. Walking and leisure-time activity and risk of hip fracture in postmenopausal women. JAMA 2002;288:2300-6.
29. Manson JE, Greenland P, LaCroix AZ, et al. Walking compared with vigorous exercise for the prevention of cardiovascular events in women. N Engl J Med 2002;347:716-25.
30. Roy DK, O'Neill TW, Finn JD, et al. Determinants of incident vertebral fracture in men and women: results from the European Prospective Osteoporosis Study (EPOS). Osteoporos Int 2003;14:19-26.

31. Yu S, Yarnell JW, Sweetnam PM, Murray L. What level of physical activity protects against premature cardiovascular death? The Caerphilly study. *Heart* 2003;89:502-6.
32. Bak H, Petersen L, Sorensen TI. Physical activity in relation to development and maintenance of obesity in men with and without juvenile onset obesity. *Int J Obes Relat Metab Disord* 2004;28:99-104.
33. Hernelahti M, Kujala U, Kaprio J. Stability and change of volume and intensity of physical activity as predictors of hypertension. *Scand J Public Health* 2004;32:303-9.
34. Hu G, Barengo NC, Tuomilehto J, Lakka TA, Nissinen A, Jousilahti P. Relationship of physical activity and body mass index to the risk of hypertension: a prospective study in Finland. *Hypertension* 2004;43:25-30.
35. Petersen L, Schnohr P, Sorensen TI. Longitudinal study of the long-term relation between physical activity and obesity in adults. *Int J Obes Relat Metab Disord* 2004;28:105-12.
36. Tammelin T, Laitinen J, Nayha S. Change in the level of physical activity from adolescence into adulthood and obesity at the age of 31 years. *Int J Obes Relat Metab Disord* 2004;28:775-82.

2. 健康の維持・増進に必要な体力（最大酸素摂取量）決定に参考とした文献

1. Blair SN, Goodyear NN, Gibbons LW, Cooper KH. Physical fitness and incidence of hypertension in healthy normotensive men and women. *Jama* 1984;252:487-90.
2. Ekelund LG, Haskell WL, Johnson JL, Whaley FS, Criqui MH, Sheps DS. Physical fitness as a predictor of cardiovascular mortality in asymptomatic North American men. The Lipid Research Clinics Mortality Follow-up Study. *N Engl J Med* 1988;319:1379-84.
3. Blair SN, Kohl HW, 3rd, Paffenbarger RS, Jr., Clark DG, Cooper KH, Gibbons LW. Physical fitness and all-cause mortality. A prospective study of healthy men and women. *Jama* 1989;262:2395-401.
4. Blair SN, Kohl HW, 3rd, Barlow CE, Gibbons LW. Physical fitness and all-cause mortality in hypertensive men. *Ann Med* 1991;23:307-12.
5. Blair SN, Kohl HW, Gordon NF, Paffenbarger RS, Jr. How much physical activity is good for health? *Annu Rev Public Health* 1992;13:99-126.
6. Hein HO, Suadicani P, Gyntelberg F. Physical fitness or physical activity as a predictor of ischaemic heart disease? A 17-year follow-up in the Copenhagen Male Study. *J Intern Med* 1992;232:471-9.
7. Kohl HW, Gordon NF, Villegas JA, Blair SN. Cardiorespiratory fitness, glycemic status, and mortality risk in men. *Diabetes Care* 1992;15:184-92.

8. Blair SN, Kohl HW, Barlow CE. Physical activity, physical fitness, and all-cause mortality in women: do women need to be active? *J Am Coll Nutr* 1993;12:368-71.
9. Sandvik L, Eriksson J, Thaulow E, Eriksson G, Mundal R, Rodahl K. Physical fitness as a predictor of mortality among healthy, middle-aged Norwegian men. *N Engl J Med* 1993;328:533-7.
10. Sawada S, Tanaka H, Funakoshi M, Shindo M, Kono S, Ishiko T. Five year prospective study on blood pressure and maximal oxygen uptake. *Clin Exp Pharmacol Physiol* 1993;20:483-7.
11. Blair SN, Kohl HW, 3rd, Barlow CE, Paffenbarger RS, Jr., Gibbons LW, Macera CA. Changes in physical fitness and all-cause mortality. A prospective study of healthy and unhealthy men. *Jama* 1995;273:1093-8.
12. Blair SN, Kampert JB, Kohl HW, 3rd, et al. Influences of cardiorespiratory fitness and other precursors on cardiovascular disease and all-cause mortality in men and women. *Jama* 1996;276:205-10.
13. Kampert JB, Blair SN, Barlow CE, Kohl HW, 3rd. Physical activity, physical fitness, and all-cause and cancer mortality: a prospective study of men and women. *Ann Epidemiol* 1996;6:452-7.
14. Lynch J, Helmrich SP, Lakka TA, et al. Moderately intense physical activities and high levels of cardiorespiratory fitness reduce the risk of non-insulin-dependent diabetes mellitus in middle-aged men. *Arch Intern Med* 1996;156:1307-14.
15. Lee CD, Jackson AS, Blair SN. US weight guidelines: is it also important to consider cardiorespiratory fitness? *Int J Obes Relat Metab Disord* 1998;22 Suppl 2:S2-7.
16. Lee CD, Blair SN, Jackson AS. Cardiorespiratory fitness, body composition, and all-cause and cardiovascular disease mortality in men. *Am J Clin Nutr* 1999;69:373-80.
17. Wei M, Gibbons LW, Mitchell TL, Kampert JB, Lee CD, Blair SN. The association between cardiorespiratory fitness and impaired fasting glucose and type 2 diabetes mellitus in men. *Ann Intern Med* 1999;130:89-96.
18. 澤田享、武藤孝司. 日本人男性における有酸素能力と生命予後に関する縦断的研究. 日本公衆衛生学雑誌 1999;46:113-121.
19. Lakka TA, Laukkanen JA, Rauramaa R, et al. Cardiorespiratory fitness and the progression of carotid atherosclerosis in middle-aged men. *Ann Intern Med* 2001;134:12-20.
20. Laukkanen JA, Lakka TA, Rauramaa R, et al. Cardiovascular fitness as a predictor of mortality in men. *Arch Intern Med* 2001;161:825-31.
21. Farrell SW, Braun L, Barlow CE, Cheng YJ, Blair SN. The relation of body mass index, cardiorespiratory fitness, and all-cause mortality in women. *Obes Res* 2002;10:417-23.

22. Myers J, Prakash M, Froelicher V, Do D, Partington S, Atwood JE. Exercise capacity and mortality among men referred for exercise testing. *N Engl J Med* 2002;346:793-801.
23. Carnethon MR, Gidding SS, Nehgme R, Sidney S, Jacobs DR, Jr., Liu K. Cardiorespiratory fitness in young adulthood and the development of cardiovascular disease risk factors. *Jama* 2003;290:3092-100.
24. Evenson KR, Stevens J, Cai J, Thomas R, Thomas O. The effect of cardiorespiratory fitness and obesity on cancer mortality in women and men. *Med Sci Sports Exerc* 2003;35:270-7.
25. Gulati M, Pandey DK, Arnsdorf MF, et al. Exercise capacity and the risk of death in women: the St James Women Take Heart Project. *Circulation* 2003;108:1554-9.
26. Kurl S, Laukkanen JA, Rauramaa R, Lakka TA, Sivenius J, Salonen JT. Cardiorespiratory fitness and the risk for stroke in men. *Arch Intern Med* 2003;163:1682-8.
27. Mora S, Redberg RF, Cui Y, et al. Ability of exercise testing to predict cardiovascular and all-cause death in asymptomatic women: a 20-year follow-up of the lipid research clinics prevalence study. *Jama* 2003;290:1600-7.
28. Sawada SS, Lee IM, Muto T, Matuszaki K, Blair SN. Cardiorespiratory fitness and the incidence of type 2 diabetes: prospective study of Japanese men. *Diabetes Care* 2003;26:2918-22.
29. Church TS, Cheng YJ, Earnest CP, et al. Exercise capacity and body composition as predictors of mortality among men with diabetes. *Diabetes Care* 2004;27:83-8.
30. Katzmarzyk PT, Church TS, Blair SN. Cardiorespiratory fitness attenuates the effects of the metabolic syndrome on all-cause and cardiovascular disease mortality in men. *Arch Intern Med* 2004;164:1092-7.
31. Stevens J, Evenson KR, Thomas O, Cai J, Thomas R. Associations of fitness and fatness with mortality in Russian and American men in the lipids research clinics study. *Int J Obes Relat Metab Disord* 2004;28:1463-70.

3. 健康の維持・増進に必要な体力（筋力、その他）決定に参考とした文献

1. Fujita Y, Nakamura Y, Hiraoka J, et al. Physical-strength tests and mortality among visitors to health-promotion centers in Japan. *J Clin Epidemiol* 1995;48:1349-59.
2. Nguyen TV, Eisman JA, Kelly PJ, Sambrook PN. Risk factors for osteoporotic fractures in elderly men. *Am J Epidemiol* 1996;144:255-63.
3. Seeley DG, Kelsey J, Jergas M, Nevitt MC. Predictors of ankle and foot fractures in older women. The Study of Osteoporotic Fractures Research Group. *J Bone Miner Res* 1996;11:1347-55.

4. Schroll M, Avlund K, Davidsen M. Predictors of five-year functional ability in a longitudinal survey of men and women aged 75 to 80. The 1914-population in Glostrup, Denmark. *Aging (Milano)* 1997;9:143-52.
5. Rantanen T, Masaki K, Foley D, Izmirlian G, White L, Guralnik JM. Grip strength changes over 27 yr in Japanese-American men. *J Appl Physiol* 1998;85:2047-53.
6. Anstey KJ, Luszcz MA, Giles LC, Andrews GR. Demographic, health, cognitive, and sensory variables as predictors of mortality in very old adults. *Psychol Aging* 2001;16:3-11.
7. Al Snih S, Markides KS, Ray L, Ostir GV, Goodwin JS. Handgrip strength and mortality in older Mexican Americans. *J Am Geriatr Soc* 2002;50:1250-6.
8. Katzmarzyk PT, Craig CL. Musculoskeletal fitness and risk of mortality. *Med Sci Sports Exerc* 2002;34:740-4.
9. Lee SH, Dargent-Molina P, Breart G. Risk factors for fractures of the proximal humerus: results from the EPIDOS prospective study. *J Bone Miner Res* 2002;17:817-25.
10. Metter EJ, Talbot LA, Schrager M, Conwit R. Skeletal muscle strength as a predictor of all-cause mortality in healthy men. *J Gerontol A Biol Sci Med Sci* 2002;57:B359-65.
11. Albrand G, Munoz F, Sornay-Rendu E, DuBoeuf F, Delmas PD. Independent predictors of all osteoporosis-related fractures in healthy postmenopausal women: the OFELY study. *Bone* 2003;32:78-85.
12. Rantanen T, Volpato S, Ferrucci L, Heikkinen E, Fried LP, Guralnik JM. Handgrip strength and cause-specific and total mortality in older disabled women: exploring the mechanism. *J Am Geriatr Soc* 2003;51:636-41.
13. Stel VS, Smit JH, Pluijm SM, Lips P. Balance and mobility performance as treatable risk factors for recurrent falling in older persons. *J Clin Epidemiol* 2003;56:659-68.
14. Metter EJ, Talbot LA, Schrager M, Conwit RA. Arm-cranking muscle power and arm isometric muscle strength are independent predictors of all-cause mortality in men. *J Appl Physiol* 2004;96:814-21.